


Ramachandran Plot

1zoy



Plot statistics

Residues in most favoured regions [A,B,L]	795	84.4%
Residues in additional allowed regions [a,b,l,p]	134	14.2%
Residues in generously allowed regions [~a,~b,~l,~p]	11	1.2%
Residues in disallowed regions	2	0.2%


Number of non-glycine and non-proline residues	942	100.0%
Number of end-residues (excl. Gly and Pro)	7	
Number of glycine residues (shown as triangles)	97	
Number of proline residues	46	

Total number of residues	1092	

Based on an analysis of 118 structures of resolution of at least 2.0 Angstroms and R-factor no greater than 20%, a good quality model would be expected to have over 90% in the most favoured regions.

Ramachandran plots for all residue types


1zoy



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0Å or better.

Ramachandran plots for all residue types


1zoy



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0Å or better.

Ramachandran plots for all residue types


1zoy



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0Å or better.

Chi1-Chi2 plots


1zoy



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0Å or better.

Chi1-Chi2 plots


1zoy



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0Å or better.

Main-chain parameters

1zoy




Plot statistics

Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. %-tage residues in A, B, L	942	84.4	78.2	10.0	0.6	Inside
b. Omega angle st dev	1088	1.2	6.0	3.0	-1.6	BETTER
c. Bad contacts / 100 residues	91	8.3	9.0	10.0	-0.1	Inside
d. Zeta angle st dev	995	1.0	3.1	1.6	-1.3	BETTER
e. H-bond energy st dev	698	0.8	0.9	0.2	-0.6	Inside
f. Overall G-factor	1092	0.3	-0.6	0.3	2.7	BETTER

Side-chain parameters

1zoy




Plot statistics

Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. Chi-1 gauche minus st dev	135	15.5	21.8	6.5	-1.0	Inside
b. Chi-1 trans st dev	308	17.2	22.0	5.3	-0.9	Inside
c. Chi-1 gauche plus st dev	417	14.5	20.5	4.9	-1.2	BETTER
d. Chi-1 pooled st dev	860	15.8	21.2	4.8	-1.1	BETTER
e. Chi-2 trans st dev	252	13.8	22.6	5.0	-1.8	BETTER


Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed

STQYPVVDHEFDVVVGAGGAGLRAAFGLSEAGFNTACVTKLFPTRSHTVAAQGGINAALGNMEEDNWRWHFYDTVKGSIDLGDQDAIHYMTEQAPASVV

f. Max. deviation (see listing)




g. G-factors




Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed

FDRGMVWNTDLVETLELQNLMLCALQTIYGAEARKE SRGAHAREDFKERVDEYDYSKPIQGQKKPFQEHWRKHTLSYVDVKTGKVSLEYRPVIDKTLNE

f. Max. deviation (see listing)




g. G-factors




Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




Residue properties

1zoy


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




Key:- Helix Beta strand Random coil Accessibility shading: Buried Accessible


e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors



Main-chain bond lengths

1zoy




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

Main-chain bond angles

1zoy




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

Main-chain bond angles

1zoy




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

RMS distances from planarity

1zoy




Histograms showing RMS distances of planar atoms from best-fit plane.
Black bars indicate large deviations from planarity: RMS dist > 0.03 for rings, and > 0.02 otherwise.

▶ signifies data points off the graph in the direction shown.

Distorted geometry


1zoy

Main-chain bond lengths




Bonds differing by $> 0.05\text{\AA}$ from small-molecule values. Values shown: "ideal", difference, actual

Main-chain bond angles



Bond angles differing by > 10.0 degrees from small-molec values. Values shown: "ideal", actual, diff.

Planar groups



Sidechains with RMS dist. from planarity $> 0.03\text{\AA}$ for rings, or $> 0.02\text{\AA}$ otherwise. Value shown is RMS dist.